

Angry? Tired? Feeling hopeless?

It's ok, it happens
to a lot of guys.



**Depression and Bipolar
Support Alliance**

We've been there. We can help.

Men and Depression

David has a great job, a nice home, and a loving family. Everyone thinks he has it made. But almost every night he comes home from a hard day's work, has a few beers, and spends the evening on the couch. He's bored with everything.

Bob blows up at the smallest things – he gets so agitated in traffic that his face turns red, he punches the steering wheel and yells at other drivers.

Jim is exhausted every night when he gets out of class. He doesn't hang out with his buddies anymore and hasn't seen his girlfriend in weeks. He just sits alone in his dorm room watching television.

These scenes are common. Who hasn't been frustrated with traffic, been tired at the end of the day or become bored with things now and then? We all have. But when feelings of irritability or sadness last for a long time, become overwhelming or start to interfere with other parts of your life, it might be a sign of something more serious: depression.

You've probably heard about depression before. People may have told you it's not a real illness and that a person should just snap out of it, or that real men don't talk about their feelings. The fact is that depression *is* a real illness just like diabetes or asthma, that affects over six million men in the United States each year. It is nothing to be embarrassed about. It happens to a lot of guys.

Clinical depression is different from normal feelings of irritability or sadness in three ways.

How intense the mood is: With depression, the feelings are much more intense.

How long it lasts: A bad mood is usually gone in a few days, but clinical depression lasts two weeks or longer.

How much it interferes with your life: A bad mood does not keep you from going to work or school or doing the things you like to do.

There is no shame in seeking help. In fact, just the opposite is true: it takes courage to admit that you need help.

You're Not Alone

Depression affects people of every age, race and economic status. While the illness affects both men and women, men typically have a different experience.

Men don't usually talk about feeling sad or lonely, instead, they are more likely to talk about being tired, having difficulty sleeping or losing interest in friends and enjoyable activities.

Men deal with depression in different ways, too. They tend to keep feelings to themselves. Many try to cope with their symptoms in unhealthy ways, including excessive use of alcohol or drugs, violent outbursts, dangerous activities or burying themselves in work.



For 20 years I got drunk every day and slept about 3 hours a night. Slowly my life got out of control. When I was working on getting sober, my

counselor suggested seeing a psychiatrist about depression. I didn't like the suggestion, but it only took a few weeks to realize that is was the right thing to do. Today, I follow my doctor's instructions, see my counselor regularly and I'm staying sober. I feel so much better.

– Art, 51 years old, married, father of two.

Symptoms of Depression

Not everyone experiences depression the same way. Some people experience many symptoms; some just a few. If you experience five or more of these symptoms for two weeks or longer, or if any of them interfere with work, school, family or social activities, talk to your doctor or a mental health professional. Do not try to diagnose or treat yourself.

Symptoms of Depression

- Irritable, empty, sad mood most of the day
- No interest in or pleasure from activities
- Major changes in appetite or body weight
- Insomnia or sleeping too much
- Feelings of restlessness
- Fatigue, exhaustion, lack of energy
- Feelings of worry pessimism or worthlessness
- Inability to concentrate or make decisions
- Thoughts of death or suicide

If you or someone you know has thoughts of death or suicide, contact a medical professional, clergy member, loved one or friend immediately or call the National Hopeline Network at (800) 442-HOPE (4673).

Bipolar Disorder

Bipolar disorder (also known as manic depression) is a treatable medical illness characterized by changes in mood, thought, energy and behavior. With bipolar disorder, a person's mood can alternate between the two poles of mania (highs) and depression (lows). Mood swings can last for days, hours, weeks or months. Unfortunately, bipolar disorder is often misdiagnosed. When you talk to your doctor, think about how you've been feeling over the past several months and tell him or her about any changes in mood you've noticed, even periods of high energy that may have felt good at the time.



I didn't realize I had depression until I was out of high school. At the age of 15, I felt 50 and was ready to die. I felt I had lived long enough and was tired of trying to fight for my life. Once I started to talk about my "secret" I realized I wasn't alone and that helped me seek treatment for my illness.

– Gregg, 20 years old

Symptoms of Mania

- Feeling energetic, high, better than good, or unusually irritable for at least one week
- Extremely high self-esteem; feeling like you can do anything
- Decreased need for sleep without feeling tired
- Talking more than usual
- Racing thoughts
- Distracted easily
- Increase in goal-directed activity, restlessness
- Excessive pursuit of pleasure (e.g., spending sprees, sexual indiscretions) without thought of consequences

Because depression and bipolar disorder are illnesses that affect moods, thoughts and behaviors, they are often referred to as mood disorders.

Treatments Work

Typically, the most effective treatment plans include three things: Talk therapy, medication and support.

Talk Therapy

Psychotherapy (talk therapy) is an important part of treatment for many people and can help you explore behaviors and modify patterns that contribute to your depression. It's important to be honest with your therapist and not be afraid to talk about things that you might think are embarrassing. Work with your therapist early in your treatment to set goals.

Medication

You and your doctor should work together to determine if medication is the right treatment option for you. The medications used to treat mood disorders are not addictive and will not alter your personality. It can take time to feel their full effect – sometimes as long as six weeks – so it's important to stick with it.

Medication Side Effects

Some people experience side effects from medication, such as dry mouth, drowsiness, restlessness, weight gain, nausea or sexual dysfunction. Often, side effects will lessen or disappear in a couple of weeks. Your doctor may suggest adjusting the medication dose or trying a different medication if your side effects continue. **Never stop taking your medication or change your dose without discussing it with your doctor.**

Herbal Remedies

Some herbal remedies and supplements have been marketed for treating depression. It is important to remember that "natural" does not always mean safe. Some herbal remedies may weaken the affects of other medications you are taking, and may affect your heart, liver or kidneys. Talk with your doctor before trying any herbal remedy.



Peer Support

Peer-to-peer support groups can provide a place to learn from other men who have gone through what you are going through. Meeting others can help you understand that your illness does not define who you are and you can learn from others who have been there.

DBSA has over 1,000 support groups across the country. Each group has a professional advisor and appointed facilitator. Members are people living with depression or bipolar disorder and their friends and family. To find the DBSA support group near you, visit our web site at www.DBSAlliance.org/findsupport.html or call (800) 826-3632.



Communicating with Your Health Professional

Open and ongoing communication with your health provider(s) is essential. You may want to take notes during your discussion, as there will be a lot of information for you to remember.

Things to tell your health provider(s)

- How you have been feeling and for how long.
- How your day-to-day life has been affected. For example, missing work, not spending time with family and friends, drug/alcohol use, etc.
- Other illness you may have, medications (including over the counter) you are taking and the names and phone numbers of other health providers you are seeing.

Questions to ask your health provider(s)

- Is medication necessary? If so, what are the benefits and drawbacks (including side effects) of different options? How much will you be taking, how many times per day? Should you take it with or without food? What should you do if you experience side effects?
- Does he or she have recommendations for a talk therapist in your community?
- Would a peer-lead support group be helpful?
- How can you reach your provider(s) in an emergency?
- Will treatment affect any other health conditions, e.g., diabetes, heart disease?
- When should you expect to start feeling better?

Wellness Lifestyle

As with any illness, maintaining a healthy lifestyle will help you manage your depression. Try to incorporate the following into your daily life.

- Get plenty of sleep.
- Exercise regularly.
- Spend time with friends and family and do things you enjoy.
- Watch your diet. Eat balanced meals. Avoid caffeine, sugar and heavily salted foods.
- Follow your treatment plan: see your talk therapist, take your medications as prescribed, and attend your support group meetings.
- Avoid drugs and alcohol.

Helping a Friend or Family Member

Men often have difficulty recognizing and talking about feelings. They may not recognize symptoms of depression in themselves or may be embarrassed to seek help. It is often a partner, family member, or a friend who points out patterns of behavior that suggest there is a problem. If you think someone you know may be struggling with depression, the best thing you can do is encourage him to talk to a health professional. Point out the changes in behavior you have noticed and suggest that he go to his doctor. Remind him that seeing a doctor for depression is no different than seeing one for high blood pressure or arthritis. Gentle reminders might be needed to get him to make the appointment. If depression is diagnosed, encourage him to stay with his treatment and provide ongoing support and feedback.



We've been there.
We can help.

Depression and Bipolar Support Alliance

The Depression and Bipolar Support Alliance (DBSA) is the leading patient-directed national organization focusing on the most prevalent mental illnesses. The organization fosters an environment of understanding about the impact and management of these life-threatening illnesses by providing up-to-date, scientifically based tools and information written in language the general public can understand. DBSA supports research to promote more timely diagnosis, develop more effective and tolerable treatments and discover a cure. The organization works to ensure that people living with mood disorders are treated equitably.

Assisted by a Scientific Advisory Board comprised of the leading researchers and clinicians in the field of mood disorders, DBSA has more than 1,000 peer-run support groups across the country. Three million people request and receive information and assistance each year. DBSA's mission is to improve the lives of people living with mood disorders.

Depression and Bipolar Support Alliance

730 N. Franklin Street, Suite 501
Chicago, Illinois 60610-7224 USA
Phone: (800) 826-3632 or (312) 642-0049
Fax: (312) 642-7243
Web site: www.DBSAAlliance.org

Visit our updated, interactive website for important information, breaking news, chapter connections, advocacy help and much more.

This brochure was reviewed by DBSA Scientific Advisory Board member George S. Alexopoulos, M.D., of the Weill Medical College of Cornell University, and Ken Heideman of DBSA Boston.

Production of this brochure was made possible by a grant from the American Psychiatric Foundation.

DBSA does not endorse or recommend the use of any specific treatments or medications for mood disorders. For advice about specific treatments or medications, individuals should consult their physicians and/or mental health professionals.